



KUNGL
TEKNISKA
HÖGSKOLAN

First Principle Study of Soft X-ray Spectroscopy


Yi Luo

Theoretical Chemistry, Department of Biotechnology
Royal Institute of Technology, Stockholm, Sweden



ALS, 2004-10-19

- Ionization/excitation of core electron (XPS-NEXAFS)
- Radiative decay of core excited/ionized state (XES-RIXS)

- core e⁻ B.Energy
- chemical environment
- satellite structures



- picture of **unoccupied** pDOS



- XES non resonantly excited
- RIXS resonantly excited selection rules
- information about **occupied** pDOS

For large molecular systems, two theoretical methods are normally used.

Hartree-Fock method (HF): without electron correlation

Wave-function based method, Ψ

Density Functional Theory (DFT): with electron correlation

Density based method: $\rho = \sum_i^N |\Psi_i|^2$

Program packages: DALTON, DeMon, GAMESS, GAUSSIAN

Density Functional Theory for X-ray Spectroscopy of Molecular Systems

Funtionals: LDA, gradient corrected, and hybrid

XPS:

Binding energy, shake-up and vibrational structures.

UPS:

Spectrum

NEXAFS:

Vibrationally resolved spectrum

XES:

**Non-resonant emission and
resonant inelastic X-ray scattering (RIXS)**

RIXS formulation

$$\begin{aligned}\langle \sigma(\omega', \omega_0) \rangle &= \sum_{vn} \frac{\omega'}{\omega} \lambda_{vn} \Phi(\omega' + \omega_{vn} - \omega_0) \\ &= \sum_{vn} \frac{\omega'}{\omega} (F\lambda_{vn}^F + G\lambda_{vn}^G + H\lambda_{vn}^H) \Phi(\omega' + \omega_{vn} - \omega_0).\end{aligned}$$

$$\lambda_{vn}^F = \sum_{\beta} F_{vn}^{\beta\beta} \sum_{\gamma} F_{vn}^{\gamma\gamma*}$$

$$F = -|e_1 e_2^*|^2 + 4|e_1 e_2|^2 - 1$$

$$\lambda_{vn}^G = \sum_{\beta\gamma} F_{vn}^{\beta\gamma} F_{vn}^{\beta\gamma*}$$



$$G = -|e_1 e_2^*|^2 - |e_1 e_2|^2 + 4$$

$$\lambda_{vn}^H = \sum_{\beta\gamma} F_{vn}^{\beta\gamma} F_{vn}^{\gamma\beta*}.$$

$$H = 4|e_1 e_2^*|^2 - |e_1 e_2|^2 - 1.$$

$$F_{vn}^{\beta\gamma} = \alpha \sum_k \omega_{vk} \omega_{nk}(v) \frac{d_{vk}^{\beta} d_{kn}^{\gamma}(v)}{\omega - \omega_{vk} + i\Gamma_{vk}}$$

Theoretical results and simulation of the experimental spectrum of metal-free Phthalocyanine H₂Pc




Benzene C1s shake-up peaks
Pyrrole C1s shake-up peaks
at about 2 eV from main line

Convoluted by gaussian curves
0.7, 0.8, 0.9 eV FWHM

First theoretical evidence of
benzene shake-up


C_{60} Calculated NEXAFS spectra

fwhm=0.5 eV




C_{70} Calculated NEXAFS spectra

fwhm=0.5 eV




Isomer Dependence



E=0

R_{cc}




$C_{2v}\text{-}C_{78}'$

Shortest: 1.373 Å
Longest: 1.477 Å
Average: 1.437 Å

E=6.34

R_{cc}




$C_{2v}\text{-}C_{78}$

Shortest: 1.366 Å
Longest: 1.476 Å
Average: 1.437 Å

E=9.48


Kcal/mol

R_{cc}




$D_3\text{-}C_{78}$

Shortest: 1.383 Å
Longest: 1.475 Å
Average: 1.437 Å



Multidimensional transition-state theory calculations for nuclear dynamics of core excited benzonitrile molecule

S. Carniato, V. Ilakovac, J.-J. Gallet, E. Kukk, and Y. Luo, *Phys. Rev. A*, in press




Carbon Nitride

Carbon nitride is today commercially used as a protective coating on hard discs and recorder heads by, e.g., IBM

New fullerene-like phase of CN_x (x = 0.1-0.3) films were discovered in 1995 in Linköping, Sweden. The phase is extremely hard and compliant (up to 95% elastic recovery!).


Many XPS and XAS studies have been reported.

The possible bonding structures have been under debate




Model I


N1s XAS






Model II




N1s XAS





SXAS N K-edge spectrum from a magnetron sputtered $\text{CN}_{0.25}$ films, grown at 350 °C (open circles), and calculated spectra from nitrile structures (N1), pyridine-like N (N2) and graphite-like N (N3). The relative intensity for the N1 spectra has been scaled to 50%, relative to N2 and N3.




Experimental (open circles) SXES N K emission spectra achieved using different excitation energies (marked by arrows in XAS), and calculated spectra (solid lines) from nitrile structures (N1), pyridine-like N (N2) and graphite-like N (N3).

Bonding Sites and Energies:

System	B.E. (eV/atom)	distance Ti-C (Å)
--------	----------------	-------------------


$C_{60}Ti$		
Bridge	3.57	2.03
5-ring	3.35	1.92(1), 2.09(2), 2.37(2)
6-ring	4.01	2.16(2), 2.17(4)

$C_{60}Ti_2$		
Bridge	3.43	2.03
5-ring	3.42	1.94(1), 2.09(2), 2.39(2)
6-ring	3.78	2.17(2), 2.21(4)





Bonding Formation

Dewar, Chatt and Duncanson (DCD) model



C_{60} Ti calculated RIXS detuning spectra







RIXS of ethylene on Cu(110)

L.Triguero, Y.Luo, L.G.M.Pettersson, H.Ågren, P.Västerlein, M.Weinelt,
A.Föhlisch, J.Hasselström, O.Karis, and A.Nilsson, *Phys.Rev.B* 59,5189 (1999)




Theoretical model: C₂H₄/Cu₈₆ + DFT

Experiment: ALS beamline 8.0





Bonding formation of Ethylen on Cu(110)



experiment




X-ray absorption of water




XES of Liquid Water





Representative water clusters



SYM




A-ASYM




D-ASYM


X-ray absorption





X-ray emission







Effects of hydrogen bond on the electronic structure of liquid water






Effects of local arrangements on the covalency of hydrogen bonding





Dynamics of core-excitation in liquid water



Normalized intensity



Liquid Methanol


- (1) Methanol crystal: infinite hydrogen-bonded chains; liquid methanol: non consensus after decades of studies
- (2) Based on neutron diffraction data: It was found experimentally that **chains up to ten molecules** with average of **six molecules** in liquid methanol [Yamaguchi, K. Hidaka, and A.K. Soper, *Molecular Physics* **96**, 1159 (1999)]; Haughney et al., *J. Phys. Chem.* **91**, 4934 (1987); S. & Joarder, R. N., *J. Chem. Phys.* **99**, 2032 (1993) revealed a different picture of **cyclic hexamers** (six-rings); Using the same diffraction data, Tanaka found **linear trimer and tetramer chains** [*Bull. Chem. Soc. Jpn.* **58**, 270 (1985)]
- (3) All molecular dynamics /Monte Carlo simulations give chain structures of different sizes.

Experimental RIXS spectra of methanol liquid




Possible structure of liquid methanol?


Chains





Rings





Six Methanol Molecules





LUMO+1




LUMO





HOMO





HOMO-1





Intensity (a.u.)



Intensity (a.u.)




Molecules in the pure liquid methanol predominantly persist as hydrogen-bonded chains and rings with six and/or eight molecules of equal abundance.



rings

Intensity (a.u.)



Experiment

Theory

10

9

8

7

6

5

4

3

240


260

280

300

Energy (eV)

chains



Experiment

Theory

10

9

8

7

6

5

4

3

240



260



280

300

Energy (eV)

XES of methanol/water mixtue








We find evidence of incomplete mixing at the microscopic level, and of water molecules bridging methanol chains to form rings.

Sheet




Folding



sideview



topview



Main experimental collaborators

Jinghua Guo	ALS
Andreas Augustsson	Uppsala University
Jan-Erik Rubensson	Uppsala University
Joseph Nordgren	Uppsala University

Main theoretical collaborator

Stepan Kashtanov	Royal Institute of Technology / ALS
Barbara Brena	Royal Institute of Technology
Hans Ågren	Royal Institute of Technology

Stephane Carniato P&M Curie University, Paris

Fundings

ALS (JingHua Guo)
Uppsala University (Joseph Nordgren)
Carl-Trygger Foundation (CTS)